

Case Report

Effect of diode laser frenectomy on tongue anatomy and function – Clinical cases



Catarina Teixeira Alves^{1,*} , Cristina Cardoso Silva^{1,2} , Otília Pereira-Lopes^{1,2} 

¹ Faculdade de Ciências da Saúde (FCS-UFP), Universidade Fernando Pessoa, Porto, Portugal

² Instituto de Investigação, Inovação e Desenvolvimento (FP-I3ID), Universidade Fernando Pessoa, Porto, Portugal

ARTICLE INFO

Article history:

Received 23 April 2024

Accepted 10 March 2025

Available online 8 May 2025

Keywords:

Ankyloglossia

Laser

Lingual frenulum

Speech

Surgery

ABSTRACT

This article describes two clinical cases of pediatric patients aged 9 years who were diagnosed with alteration of the lingual frenulum and subsequently underwent diode laser frenectomy after multidisciplinary clinical evaluation. Post-surgical functional and phonetic changes were evaluated using the Marchesan Protocol for Lingual Frenulum Assessment and the Wong-Baker Pain Face Scale. The cases presented showed improvements in anatomical parameters and tongue motricity after frenectomy. However, analysis of phonetic changes revealed that immediate improvements were most evident in the less complex sounds, while more challenging phonetic changes (such as reservation, omission, and substitutions and phonemes) required further rehabilitation. This result reinforces that surgical intervention is only one component of the therapeutic process and must be complemented by an interdisciplinary approach that includes speech therapy adapted to the patient's specific needs to improve clinical and functional results. (Rev Port Estomatol Med Dent Cir Maxilofac. 2025;66(x):xxx-xxx)

© 2025 Sociedade Portuguesa de Estomatologia e Medicina Dentária.

Published by SPEDM. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Ankyloglossia (from the Greek “ankylo,” meaning “rigid,” and “glossa,” meaning “tongue”) is a partial or complete congenital anomaly of the lingual frenulum, where it is abnormally short or inserted too close to the lingual apex.¹ According to the International Affiliation of Tongue-Tie Professionals (IATP), this condition is also known as “symptomatic tongue-tie” or “symptomatic ankyloglossia.” It can be classified as anterior or posterior. In anterior ankyloglossia, the tongue's

protrusion movements are restricted and originate a “heart-shaped” appearance due to the involvement of the tip of the tongue. Posterior ankyloglossia is difficult to identify and, unlike the former, does not involve the tip of the tongue but rather the thickening of the frenulum, which shows a fibrous appearance.²

Ankyloglossia's prevalence varies from 0.1% to 10.7%,³ and its incidence from 0.02% to 5%.⁴⁻⁶ Clinically, it results in restriction of normal lingual movements and can be associated with functional limitations, such as interference with feeding, suck-

* Corresponding author.

E-mail address: catarinatexeira@hotmail.com (Catarina Teixeira Alves).

<http://doi.org/10.24873/j.rpemd.2025.04.1424>

1646-2890/© 2025 Sociedade Portuguesa de Estomatologia e Medicina Dentária. Published by SPEDM.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ing, and swallowing, and phonetic disorders with speech imprecision due to exchange or distortion of lingual sounds, among others.⁷ These phonetic and functional challenges underline the importance of a multidisciplinary approach, where dentistry and speech therapy intervention play key roles. While the speech therapist addresses the functional rehabilitation of speech and swallowing, the dentist focuses on the structural and surgical correction of the lingual frenulum. This collaboration is particularly critical in pediatric cases, where early intervention can have a significant impact on developmental outcomes.

There are three main surgical options to correct the altered lingual frenulum: frenotomy, frenectomy, and frenuloplasty. In summary, frenotomy is most commonly used in neonates and infants due to its simplicity and effectiveness, while frenectomy and frenuloplasty are the procedures of choice for corrective therapy in children, adolescents, and adults.⁸

The most common technique to correct tongue-tie is using a conventional scalpel. However, several clinicians have suggested using a soft-tissue laser as a favorable technique for correcting short lingual and/or labial frenulae.^{9,10} Laser technology can provide a more precise incision in a shorter operative time, cauterize small blood and lymphatic vessels through its hemostatic and anti-inflammatory effects, minimize discomfort and pain during the intraoperative period, significantly reduce postoperative discomfort, and facilitate the healing process of the surgical wound.¹¹ Moreover, numerous studies have documented that the incision in the soft tissues of the oral cavity with a laser is less painful than the incision with a scalpel and that patients treated with a laser have less pain and fewer functional complications after surgery than those who underwent conventional surgery. However, no specific surgical approach has been shown to be superior to the others to the point of being recommended as a gold standard.¹²

Ideally, this condition should be diagnosed shortly after birth. It may also be diagnosed during the baby's first dental visit, which, according to the European Academy of Paediatric Dentistry (EAPD) and the American Academy of Pediatric Dentistry (AAPD), should be at the time of eruption of the first primary tooth or one year of age, if no teeth have erupted. Numerous diagnostic instruments are available to the dentist for lingual frenulum anomalies, the most notable of which is the Marchesan's Lingual Frenulum Assessment Protocol (hereafter, Marchesan Protocol),¹³⁻¹⁵ whose primary objective is to evaluate and score the various lingual frenular characteristics, normal or abnormal.

The integration of dentistry and speech therapy is crucial in managing tongue-tie cases, as these professionals address different but complementary aspects of the condition. While dentists provide surgical solutions, speech therapists ensure proper functional rehabilitation through exercises to optimize oral-motor skills. This multidisciplinary approach is essential to achieving structural correction and long-term functional improvement likewise, particularly in children whose speech and feeding skills are still developing.

We present two clinical cases to measure and compare the functional and/or phonetic activity before and after the surgical treatment of altered lingual frenula in two chil-

dren. Our aim was to monitor the postoperative condition after laser frenectomy and quantify the patients' discomfort by understanding the substantial differences felt in their functional and/or phonetic activity after the therapeutic approach.

Clinical Cases

Two 9-year-old pediatric patients, named Clinical Case 1 (C1) and Clinical Case 2 (C2), were evaluated at the Pedagogical Clinic of Dental Medicine at Fernando Pessoa University for difficulties in speech and tongue mobility due to short lingual frenula. Both patients had no previous surgical intervention and were undergoing regular speech therapy.

The Marchesan Protocol was used to evaluate the lingual frenulum at three different time points: preoperatively and postoperatively at 1 and 4 weeks. Photographs were taken for comparative analysis. Frenectomy with the Lasotronic Smart M® diode laser (Lasotronic, Poland) at 980 nm was the procedure chosen after obtaining informed consent from the parents.

In C1, no systemic pathology was reported. The patient had been bottle-fed for up to 2 years and had onychophagia. Clinical evaluation revealed significant phonetic changes without affecting chewing or swallowing and confirmed anterior ankyloglossia with a short and anteriorized frenulum (Figure 1). Before the procedure, local anesthesia was performed using 2% lidocaine with epinephrine (1:100000), applied through infiltration. The procedure lasted 15 minutes.

In C2, the patient had a history of atopic skin and had been breastfed and then bottle-fed. There was a family history of ankyloglossia. Clinical evaluation indicated posterior ankyloglossia with a fibrous frenulum (Figure 2). Similar to C1, phonetic changes were noted without affecting other oral functions. Before the procedure, local anesthesia with 2% lidocaine and epinephrine (1:100000) was used. The procedure lasted 15 minutes.



Figure 1. Initial situation of patient C1 – maximum lingual elevation achieved with the mouth open.



Figure 2. Initial situation of patient C2 – maximum lingual elevation achieved with the mouth open.

Laser frenectomy was performed in both cases, using local anesthesia and lingual stabilization techniques for surgical precision. The laser was adjusted to specifications that minimize discomfort and maximize effectiveness. Postoperative medication included paracetamol for pain control, and no additional medications were necessary.

Preoperative analysis using the Marchesan Protocol revealed an abnormal lingual frenulum with scores of 6 for C1 and 4 for C2. A score of 6 in the Marchesan Protocol indicates a severe limitation in tongue mobility, while a score of 4 represents a moderate limitation. Postoperative evaluations showed improvement, with C1 maintaining an altered frenulum (score 3) and C2 being completely corrected (score 2). A score of 3 reflects mild restriction of tongue movement, and a score of 2 indicates normal mobility.

Postoperative care included the administration of paracetamol and detailed instructions for pain control and oral hygiene. Follow-up consultations revealed progress in healing and tongue mobility, using the Marchesan Protocol and the Wong-Baker FACES pain rating scale to monitor patients' well-being (Tables 1, 2, and 3).

Table 1. Description of the preoperative general tests and the postoperative general test after 1 week and after 4 weeks.

General tests	Preoperative		Postoperative			
			1 week		4 weeks	
	C1	C2	C1	C2	C1	C2
Score – Changes during tongue elevation	Heart-shaped lingual apex	No alterations of the lingual apex upon tongue elevation	Heart-shaped lingual apex	No alterations of the lingual apex upon tongue elevation	Heart-shaped lingual apex	No alterations of the lingual apex upon tongue elevation
MR = 0; PR = 2	(1)	(0)	(1)	(0)	(1)	(0)
Score – Relationship between AMB and AMB with the lingual apex on the IP (%)	18.18	10.34	36.36	27.8	38.64	32.76
$\geq 50.1\% = 0$; $\leq 50\% = 1$	(1)	(1)	(1)	(1)	(1)	(1)
Score – Fixation of the FL on the floor of the oral cavity	Visible already from the inferior alveolar ridge	Visible already from the inferior alveolar ridge	Visible already from the inferior alveolar ridge	Visible already from the inferior alveolar ridge	Visible already from the inferior alveolar ridge	Visible already from the inferior alveolar ridge
Fixation of the FL on the underside of the tongue	Between the middle part and the apex	In the middle part	In the middle part	In the middle part	In the middle part	In the middle part
MR = 0; PR = 3	(2)	(1)	(1)	(1)	(1)	(1)
Score – FL clinical classification	FL changed; short and anterior	FL changed; short	Normal FL	Normal FL	Normal FL	Normal FL
MR = 0; PR = 2	(2)	(2)	(0)	(0)	(0)	(0)
Total score	6	4	3	2	3	2
MR = 0; PR = 8	($\geq 3 \rightarrow$ FL anomalous)	($\geq 3 \rightarrow$ FL altered)	($\geq 3 \rightarrow$ FL altered)	(< 3 \rightarrow FL not changed)	($\geq 3 \rightarrow$ FL altered)	(< 3 \rightarrow FL not changed)







C1 – Clinical Case 1; C2 – Clinical Case 2; AMB – maximum mouth opening; PI – incisive papilla; FL – lingual frenulum; % – percentage; mm – millimeters; MR – best result; PR – worst result.

Table 2. Description of functional tests conducted preoperatively and postoperatively after 1 week and 4 weeks.

Functional tests	Preoperative		Postoperative			
			1 week		4 weeks	
	C1	C2	C1	C2	C1	C2
Score – language mobility MR = 0; PR = 14	(4)	(4)	(2)	(2)	(0)	(2)
Score– tongue at rest MR = 0; PR = 4	On the floor of the oral cavity (1)	On the floor of the oral cavity (1)	On the floor of the oral cavity (1)	On the floor of the oral cavity (1)	On the floor of the oral cavity (1)	On the floor of the oral cavity (1)
Score – speech MR = 0; PR = 12	(9)	(12)	(0)	(9)	(0)	(9)
Score – other aspects MR = 0; PR = 10	(3)	(3)	(0)	(2)	(0)	(2)
Altered sounds	[g],[z],[j]	[t],[s],[g],[z],[r],[ch],[j]	No relevant changes to point out	[s],[z],[ch]	No relevant changes to point out	[s],[z],[ch]
Total score MR = 0; PR = 40	17 (< 25→ There may or may not be FL interference)	20 (< 25→ There may or may not be FL interference)	3	14	1	14

C1 – Clinical Case 1; C2 – Clinical Case 2; FL – lingual frenulum; MR – best result; PR – worst result.

Table 3. Description of the results of the Wong-Baker FACES pain rating scale concerning the preoperative period and the postoperative periods after 1 week and 4 weeks.

	C1 Preoperative	C1 Postoperative (1 week)	C1 Postoperative (4 weeks)	C2 Preoperative	C2 Postoperative (1 week)	C2 Postoperative (4 weeks)
Wong-Baker FACESpain rating scale – selected face						
	0	6	0	0	2	0
	No Hurt	Hurts Even More	No Hurt	No Hurt	Hurts Little Bit	No Hurt

Re-evaluations confirmed significant functional and anatomical improvements, validating the intervention performed. Both patients showed progressive recovery with a reduction in the lingual limitations initially observed (Figures 3, 4, and 5).

Discussion and Conclusions

The effect on tongue anatomy and function after diode laser frenectomy in the two clinical cases presented, C1 and C2, resulted in improvements. The use of the Marchesan Protocol proved essential for the systematical and reliable measure-

ment of changes in tongue anatomy and function after diode laser frenectomy. This finding reinforces the validity of this protocol and highlights its clinical relevance in understanding the results obtained.

Regarding the postoperative morphological and anatomical parameters of the Marchesan Protocol, the lingual frenulum was still classified as altered in C1, while it was classified as unchanged in C2. Given the persistence of lingual frenulum alterations in C1, it may be pertinent to consider what benefit could have been achieved by a deeper intervention, namely, the dissection of the genioglossus muscle.



Figure 3. 1-week post-operative control of patient C1 – maximum lingual elevation achieved with the mouth open.



Figure 5. 1-week post-operative control of patient C2 – clinical appearance of the fibrin clot formed in the surgical site treated.



Figure 4. 4-week post-operative control of patient C1 – maximum tongue elevation achieved with the mouth open.

Postoperative evaluations revealed no lingual frenulum change in C2. Despite expectations that this patient would exhibit the greatest phonetic potential improvement, C1 showed the most functional gains, with complete correction of the preoperative phonetic changes. Conversely, C2 continued to experience difficulty with correctly pronouncing some isolated and combined phonemes, such as [s], [z], [ch], and the persistence of possible compensatory mechanisms (such as mandibular deviation) was noted.

The dichotomy and apparent lack of agreement between the anatomic-morphological and functional parameters observed in these clinical cases raise doubts about the strictly linear relationship between lingual frenulum morphology and phonetic adequacy. It also questions the contributing factors, besides the frenulum anatomy, capable of influencing speech and other functional activities of the stomatognathic system,

confirming what has already been reported by other authors.¹⁶

The tailored approach to C1 and C2 highlights the importance of customizing surgical techniques to individual needs. For instance, in C1, where some alteration of the lingual frenulum persisted postoperatively, a more invasive surgical option, such as dissection of the genioglossus muscle, could have been considered to achieve better anatomical and functional outcomes. Both cases benefited from diode laser frenectomy, which ensured precise cutting with minimal discomfort, but postoperative care, including pain management with paracetamol and oral hygiene guidance, was critical to optimizing recovery. Adjunctive surgical techniques, such as lingual stabilization during the procedure, enhanced precision, and minimized postoperative complications, were also crucial. These elements underscore the importance of a comprehensive and individualized approach to clinical intervention in managing ankyloglossia.

Nevertheless, multidisciplinary involvement between the dentist and the speech therapist is highly advisable to alleviate functional and phonetic consequences that persist after surgery, as has been so widely advocated in the most recent literature on the subject.^{17,18} The speech therapist's intervention is a fundamental part of the treatment process. Speech therapy, both preoperatively and postoperatively, plays a key role in addressing compensatory mechanisms, improving articulation, and optimizing the functional outcomes of surgical intervention. In these cases, the integration of speech therapy could have further supported C2 in overcoming phonetic difficulties and reducing compensatory mandibular deviations observed postoperatively. The collaboration between dentists and speech therapists is crucial to achieving both anatomical correction and functional rehabilitation, ensuring comprehensive care for patients with ankyloglossia.

Diode laser frenectomy ensured a cleaner, drier operative field with better visualization, as the hemostatic cut was immediate and limited to the site of contact between the optical fiber and the most superficial blood vessels. The diode laser has increasingly demonstrated its potential as an ideal alter-

native to conventional frenectomy.^{9,10} Additionally, the literature has described other advantages of the diode laser when compared to the conventional or electric scalpel, such as the following: superior precision; lower risk of inducing unwanted damage to adjacent tissues; significantly reduced operation duration due to not needing suture and sterilization of the surgical site; minimization of bacterial proliferation; less painful intra and postoperative periods; better acceptance by pediatric dentistry patients; absence of postincisional adhesions, thus not compromising lingual mobility in the immediate postoperative period.¹¹

It should be noted that although the risk of complications resulting from this surgical procedure is reduced, it is not non-existent; i.e., the patient may feel slight edema on the oral cavity floor and moderate pain in the region. The most serious risks are injury to Wharton's ducts, damage to the anastomoses of the branches of the lingual nerves, and even damage to the deep lingual vein, which can cause profuse venous hemorrhage.¹⁹⁻²¹

Since the fixation site and the histological composition of the lingual frenulum cannot be modified by mobilization exercises, surgical intervention should be considered a fundamental part of the treatment plan. However, integrating speech therapy into the treatment plan is equally fundamental, as it complements surgical intervention by enhancing functional outcomes and addressing compensatory mechanisms that surgery alone cannot resolve. This multidisciplinary approach ensures a more effective and holistic resolution of the challenges posed by ankyloglossia.

In the clinical cases presented, lingual anatomy and motricity aspects improved after lingual frenectomy. However, no improvements were observed in the more complex phonetic changes, such as distortion, omission, and replacement of phonemes associated with an altered lingual frenulum. Thus, coordinated articulation with speech therapy for functional re-education becomes essential.

Conflict of interest

The authors have no conflicts of interest to declare.

Ethical disclosures

Protection of human and animal subjects. The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Confidentiality of data. The authors declare that they have followed their work center protocols on access to patient data and for its publication.

Right to privacy and informed consent. The authors have obtained the written informed consent of the patients or subjects mentioned in the article. The corresponding author is in possession of this document.

CREDIT AUTHORSHIP CONTRIBUTION STATEMENT

Catarina Teixeira Alves: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft. **Cristina Cardoso Silva:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing – review & editing. **Otilia Pereira-Lopes:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing – review & editing.

ORCID

Catarina Teixeira Alves  0009-0006-2091-5752

Cristina Cardoso Silva  0000-0003-3757-596X

Otilia Pereira-Lopes  0000-0002-6199-4091

REFERENCES

1. Fujinaga CI, Chaves JC, Karkow IK, Klossowski DG, Silva FR, Rodrigues AH. Lingual frenum and breast feeding: descriptive study. *Audiol Commun Res.* 2017;22:e1762.
2. Arruda EMG, Campos F, Vasconcelos RG, Vasconcelos MG. Repercussão da anquiloglossia em neonatos: diagnóstico, classificação, consequências clínicas e tratamento. *Rev Salusvita (Bauru).* 2019;38:1107-26.
3. Suter VGA, Bornstein MM. Ankyloglossia: facts and myths in diagnosis and treatment. *J Periodontol.* 2009;80:1204-19.
4. Kupietzky A, Botzer E. Ankyloglossia in the infant and young child: clinical suggestions for diagnosis and management. *Pediatr Dent.* 2005;27:40-6.
5. Buryk M, Bloom D, Shope T. Efficacy of neonatal release of ankyloglossia: a randomized trial. *Pediatrics* 2011;128:280-8.
6. Edmunds J, Miles SC, Fulbrook P. Tongue-tie and breastfeeding: a review of the literature. *Breastfeed Rev.* 2011;19:19-26.
7. Braga LAS, Silva J, Pantuzzo CL, Motta AR. Prevalência de alteração no frênulo lingual e suas implicações na fala de escolares. *Ver. CEFAC.* 2009;11(suppl 3):378-90.
8. Tuli A, Singh A. Monopolar diathermy is used for the correction of ankyloglossia. *J Indian Soc Pedod Prev Dent.* 2010;28:130-3.
9. Fioravanti M, Zara F, Vozza I, Polimeni A, Sfasciotti GL. The efficacy of lingual laser frenectomy in pediatric OSAS: A randomized double-blinded and controlled clinical study. *Int J Environ Res Public Health.* 2021;18:6112.
10. Nunes JEP, Silva AL, Nunes GP, Fagundes CF, Souza MR, Navarro RS. Lasers aplicados à frenectomia em pacientes pediátricos com anquiloglossia. *Research, Society and Development.* 2021;10:e19210210434.
11. Melo AJB, Santos GM, Júnior MBS, Mendes VCO, Melo PHB. Lasers de alta potência na frenectomia, seus benefícios e limitações: revisão integrativa. *Research, Society and Development.* 2022;11:e506111234806.
12. Lima CB, Maranhão VF, Botelho KVG, Junior VES. Avaliação da anquiloglossia em neonatos por meio do teste da linguinha: um estudo de prevalência. *RFO (Passo Fundo).* 2018;22:294-7.
13. Marchesan IQ. Lingual frenulum: quantitative evaluation porposol. *Int J Orofacial Myology.* 2005;31:39-48.
14. Marchesan IQ. Protocolo de avaliação do frênulo da língua. *Rev CEFAC.* 2010;12:977-89.

15. Marchesan IQ. Lingual frenulum: classification and speech interference. *Int J Orofacial Myology*. 2004;30:31-8.
16. Pereira TS, Oliveira F, Cardoso MCAF. Association between harmful oral habits and the structures and functions of the stomatognathic system: perception of parents/guardians. *CoDAS*; 2017;29,e20150301.
17. Gomes E, Araújo FBD, Rodrigues JÁ. Freio lingual: abordagem clínica interdisciplinar da Fonoaudiologia e Odontopediatria. *Revista da Associação Paulista de Cirurgões Dentistas*. 2015;69:20-4.
18. Miranda PP, Cardoso CL, Gomes E. Interventions in the Alteration on Lingual Frenum: Systematic Review. *Int Arch Otorhinolaryngol*. 2016;20:275-80.
19. Hall DMB, Renfrew MJ. Tongue tie. *Arch Dis Child*. 2005;90:1211-5.
20. Lalakea ML, Messner AH. *Frenotomy And Frenuloplasty: If, When, And How*. Operative Techniques in Otolaryngology – Head and Neck Surgery. 2002;13:93-7.
21. Yang H, WooY, Won S, Kim D, Hu K, Kin H.. *Course and Distribution of the Lingual Nerve in the Ventral Tongue Region: Anatomical Considerations for Frenectomy*. *J Craniofac Surg*. 2009;20:1359-63.

Efeito na anatomia e função da língua após frenectomia a laser diodo – Casos clínicos

R E S U M O

O presente artigo descreve dois casos clínicos de pacientes com 9 anos de idade, diagnosticados com alteração do freio lingual e subsequentemente submetidos a frenectomia a laser diodo após avaliação clínica multidisciplinar. Para avaliar as alterações funcionais e fonéticas pós-cirúrgicas foi utilizado o Protocolo de Marchesan para Avaliação do Freio Lingual juntamente com a Escala de Faces de Dor de Wong-Baker. Nos casos apresentados, verificaram-se melhorias nos parâmetros anatômicos e de motricidade lingual após a cirurgia. Contudo, a análise das alterações fonéticas demonstrou que as melhorias imediatas foram mais evidentes nos sons menos complexos, enquanto as alterações fonéticas mais desafiantes (como distorção, omissão e substituição e fonemas) exigem reabilitação adicional. Este resultado reforça que a intervenção cirúrgica é apenas um componente do processo terapêutico e deve ser complementada por uma abordagem interdisciplinar que inclua terapia da fala adaptada às necessidades específicas do paciente, para otimizar os resultados clínicos e funcionais. (Rev Port Estomatol Med Dent Cir Maxilofac. 2025;66(x):xxx-xxx)

© 2025 Sociedade Portuguesa de Estomatologia e Medicina Dentária.

Publicado por SPEMD. Este é um artigo Open Access sob uma licença CC BY-NC-ND

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palavras-chave:

Anquiloglossia
Laser
Freio lingual
Fala
Cirurgia